

	Diseases Preventable	Schedule age of 1st dose	Number of doses
Rota Virus	Diarrhea	2 months	<ul style="list-style-type: none"> The first dose before 15 weeks of age and the last dose before 6 months of age Rotarix: 2 doses 2 months apart Rotateq: 3 doses 2 months apart
Pneumococcal	Pneumonia, meningitis, ear and sinus infection	2 months	<ul style="list-style-type: none"> Synflorix & Prevnar-13: 3 doses 2 months apart with or without booster at 15-18 months of age If Initiating vaccination after 12 months of age; <ul style="list-style-type: none"> Synflorix: between 1 to 5 years of age, 2 doses 2 months apart Prevnar-13: if older than 1 year but less than 2 years old, 2 doses 2 months apart. If older than 2 years, 1 dose
Haemophilus Influenzae Type B (Hib Booster)	Pneumonia, meningitis, sinus and ear infection	12-15 months	<ul style="list-style-type: none"> 1 booster dose Especially for those at high risk of complications such as those with: <ul style="list-style-type: none"> Anatomical /functional asplenia - thalassemia major Immunodeficiency, etc
Influenza	Flu, pneumonia, heart problems, ear and sinus infection etc	6 months of age and above	<ul style="list-style-type: none"> If younger than 9 years of age, 2 doses 4 weeks apart in the first year of vaccination, followed by 1 dose every year If older than 9 years of age or an adult, 1 dose every year <p>Annual vaccination is recommended for health care workers and especially for those at high risk of influenza related complications including people who</p> <ul style="list-style-type: none"> are pregnant are thalassemia majors have chronic heart, lung, liver, kidney disease are younger than 5 years or older than 65 years are immunocompromised have diabetes mellitus have asthma
Chicken pox	Chicken pox	Older than 1 year	<ul style="list-style-type: none"> 2 doses Routine: 1st dose at 12-15 months followed by 2nd dose at 4-6 years For children between 16 months and 13 years, 2 doses at least 3 months apart If older than 13 years of age and not vaccinated/never had chicken pox disease; 2 doses 1 month apart. Especially recommended for health care workers and childhood educators/care takers
Hepatitis A	Hepatitis	Older than 1 year	<ul style="list-style-type: none"> 2 doses, 6 months apart <p>(Note: if older than 19 years, requires adult formulation)</p>